

Prevalence of Selected Maternal and Child Health Indicators, Mississippi, Pregnancy Risk Assessment Monitoring System (PRAMS), 2021

Health Indicator	N*	%[†] (95% CI)[‡]
Nutrition		
Multivitamin use \geq 4 days a week during the month before pregnancy	883	34.5 (30.6-38.4)
Pre-pregnancy Weight		
Underweight (BMI < 18.5)	853	2.0 (0.9-3.0)
Overweight (BMI 25.0-29.9)	853	25.7 (22.1-29.4)
Obese (BMI \geq 30.0)	853	37.3 (33.3-41.3)
Substance Use		
Any cigarette smoking during the 3 months before pregnancy	865	15.7 (12.7-18.7)
Any cigarette smoking during the last 3 months of pregnancy	864	8.3 (6.0-10.5)
Any cigarette smoking postpartum	865	11.1 (8.5-13.6)
Any e-cigarette use during the 3 months before pregnancy	863	8.4 (6.4-11.1)
Any e-cigarette use during the last 3 months of pregnancy	866	3.0 (1.9-4.8)
Hookah use in the last 2 years	859	4.1 (2.4-5.5)
Any alcohol use during the 3 months before pregnancy	859	48.6 (44.5-52.8)
Intimate Partner Physical Abuse		
Experienced intimate partner violence (IPV) during the 12 months before pregnancy by a husband or partner	853	2.8 (1.5-4.2)
Experienced IPV during the 12 months before pregnancy by an ex-husband or ex-partner	853	2.4 (1.1-3.7)
Experienced IPV during pregnancy by a husband or partner / an ex-husband or ex-partner	856	3.1 (1.6-4.5)
Depression		
Self-reported depression in the 3 months before pregnancy	883	19.0 (15.8-22.3)
Self-reported depression during pregnancy	855	18.8 (15.6-22.1)
Self-reported postpartum depressive symptoms	837	20.8 (17.4-24.2)
Health Care Services		
Health care visit in the 12 months before pregnancy	877	57.8 (53.8-61.9)
Began prenatal care in 1st trimester	851	85.5 (82.6-88.3)
Had flu shot in 12 months before delivery	874	39.4 (35.4-43.5)
Had maternal postpartum checkup	841	88.1 (85.4-90.7)

Family Planning		
Mistimed	865	23.9 (20.4-27.4)
Unwanted pregnancy	865	9.2 (6.9-11.4)
Unsure about pregnancy	865	19.8 (16.6-23.1)
Intended pregnancy	865	47.1 (43.0-51.3)
Use of any postpartum contraception	845	74.8 (71.2-78.4)
Use of a long-acting reversible contraceptive method (Intrauterine Device [IUD] or contraceptive implant) postpartum	840	12.6 (9.8-15.5)
Oral Health		
Teeth cleaned during pregnancy	875	35.3 (31.4-39.3)
Pre-pregnancy Health Insurance[‡] Status		
Private insurance	864	52.7 (48.6-56.9)
Medicaid	864	25.9 (22.3-29.5)
No insurance	864	20.9 (17.5-24.3)
Health Insurance[‡] Status for Prenatal Care During Pregnancy		
Private insurance	837	44.0 (39.8-48.2)
Medicaid	837	54.6 (50.4-58.8)
No insurance	837	1.4 (0.7-2.9)
Health Insurance[‡] Status Postpartum		
Private insurance	866	41.6 (37.5-45.7)
Medicaid	866	48.6 (44.4-52.7)
No insurance	866	9.6 (7.1-12.1)
Infant Sleep Practices		
Baby most often laid on back to sleep	819	70.7 (66.9-74.5)
Breastfeeding Practices		
Ever breastfed	828	75.8 (72.2-79.4)
Any breastfeeding at 8 weeks	815	46.5 (42.3-50.7)

* Unweighted sample size

† Weighted percent

‡ 95% Confidence Interval

‡ Insurance is coded as Medicaid (Medicaid or state-named Medicaid program); Private (Private only, both Medicaid & private, any other insurance in combination with private, TRICARE or other military insurance); and No insurance [no insurance or Indian Health Service (IHS) only]