

SELECTED DEATH STATISTICS, RESIDENTS OF MISSISSIPPI, 2003

| LEADING CAUSES OF DEATH LISTED IN DESCENDING ORDER | -----NUMBER----- | | | ---MORTALITY RATE*--- | | |
|---|------------------|--------|---------------|-----------------------|-------|---------------|
| | TOTAL | WHITE | NON- WHITE | TOTAL | WHITE | NON- WHITE |
| TOTAL OF ALL CAUSES | 28,333 | 19,013 | 9,320 | 9.8 | 10.8 | 8.3 |
| HEART DISEASES | 8,662 | 5,932 | 2,730 | 300.6 | 336.4 | 244.2 |
| MALIGNANT NEOPLASMS | 5,924 | 4,006 | 1,918 | 205.6 | 227.1 | 171.6 |
| CEREBROVASCULAR DISEASE | 1,723 | 1,110 | 613 | 59.8 | 62.9 | 54.8 |
| ACCIDENTS | 1,647 | 1,147 | 500 | 57.2 | 65.0 | 44.7 |
| EMPHYSEMA AND OTHER CHRONIC LOWER RESPIRATORY DISEASES | 1,385 | 1,182 | 203 | 48.1 | 67.0 | 18.2 |
| INFLUENZA AND PNEUMONIA | 754 | 553 | 201 | 26.2 | 31.4 | 18.0 |
| NEPHRITIS, NEPHROTIC SYND, NEPHROSIS | 675 | 361 | 314 | 23.4 | 20.5 | 28.1 |
| DIABETES MELLITUS | 671 | 355 | 316 | 23.3 | 20.1 | 28.3 |
| ALZHEIMERS DISEASE | 575 | 453 | 122 | 20.0 | 25.7 | 10.9 |
| SEPTICEMIA | 537 | 313 | 224 | 18.6 | 17.7 | 20.0 |
| HYPERTENSION/HYPERTENSIVE RENAL DZ | 369 | 177 | 192 | 12.8 | 10.0 | 17.2 |
| SUICIDE | 336 | 283 | 53 | 11.7 | 16.0 | 4.7 |
| HOMICIDE | 304 | 83 | 221 | 10.6 | 4.7 | 19.8 |
| CHRONIC LIVER DISEASE AND CIRRHOSIS | 220 | 167 | 53 | 7.6 | 9.5 | 4.7 |
| PNEUMONITIS DUE TO SOLIDS & LIQUIDS | 215 | 154 | 61 | 7.5 | 8.7 | 5.4 |
| CONDITIONS ORIG IN PERINATAL PERIOD | 204 | 65 | 139 | 4.8 | 2.8 | 7.2 |
| HIV DISEASE | 188 | 33 | 155 | 6.5 | 1.9 | 13.9 |
| AORTIC ANEURYSM AND DISSECTION | 147 | 116 | 31 | 5.1 | 6.6 | 2.8 |
| PARKINSONS DISEASE | 125 | 111 | 14 | 4.3 | 6.3 | 1.2 |
| CONGENITAL MALFORMATIONS AND CHROMOSOMAL ABNORMALITIES | 114 | 53 | 61 | 4.0 | 3.0 | 5.4 |
| IN SITU, BENIGN & UNSPECIFIED NEOPLASMS | 94 | 64 | 30 | 3.3 | 3.6 | 2.7 |
| COMPLICATIONS MEDICAL/SURGICAL CARE | 63 | 31 | 32 | 2.2 | 1.8 | 2.9 |
| ANEMIAS | 57 | 25 | 32 | 2.0 | 1.4 | 2.9 |
| VIRAL HEPATITIS | 56 | 48 | 8 | 1.9 | 2.7 | 0.7 |
| ATHEROSCLEROSIS | 55 | 44 | 11 | 1.9 | 2.5 | 1.0 |
| MALNUTRITION/NUTRITIONAL DEFICIENCY | 41 | 26 | 15 | 1.4 | 1.5 | 1.3 |
| PEPTIC ULCER | 31 | 20 | 11 | 1.1 | 1.1 | 1.0 |
| CHOLELITHIASIS/GALLBLADDER DISORDER | 30 | 22 | 8 | 1.0 | 1.2 | 0.7 |
| PNEUMOCONIOSES & CHEMICAL EFFECTS | 17 | 16 | 1 | 0.6 | 0.9 | 0.1 |
| MENINGITIS | 11 | 3 | 8 | 0.4 | 0.2 | 0.7 |
| HERNIA | 11 | 5 | 6 | 0.4 | 0.3 | 0.5 |
| MATERNAL CAUSES | 7 | 3 | 4 | 16.5 | 13.0 | 20.8 |
| DISEASES OF APPENDIX | 6 | 5 | 1 | 0.2 | 0.3 | 0.1 |
| ACUTE BRONCHITIS AND BRONCHIOLITIS | 5 | 4 | 1 | 0.2 | 0.2 | 0.1 |
| INFECTIONS OF KIDNEY | 5 | 4 | 1 | 0.2 | 0.2 | 0.1 |
| TUBERCULOSIS | 4 | 4 | 0 | 0.1 | 0.2 | - |
| FEMALE PELVIC INFLAMMATORY DISEASE | 4 | 1 | 3 | 0.1 | # | 0.3 |
| LEGAL INTERVENTION | 3 | 1 | 2 | 0.1 | # | 0.2 |
| VIRAL ENCEPHALITIS | 2 | 2 | 0 | 0.1 | 0.1 | - |
| SALMONELLA INFECTION | 1 | 1 | 0 | # | # | - |
| MENINGOCOCCAL INFECTION | 1 | 1 | 0 | # | # | - |
| SYPHILIS | 1 | 0 | 1 | # | - | 0.1 |
| HYPERPLASIA OF PROSTATE | 1 | 1 | 0 | # | # | - |
| OPERATIONS OF WAR & THEIR SEQUELAE | 1 | 1 | 0 | # | # | - |
| --- NOT INCLUDED IN RANKING --- | | | | | | |
| SYMPTOMS, SIGNS, ILL-DEFINED | 431 | 258 | 173 | 15.0 | 14.6 | 15.5 |
| ALL OTHER CAUSES | 2,620 | 1,769 | 851 | 90.9 | 100.3 | 76.1 |

* TOTAL DEATHS PER 1,000 POPULATION. DEATHS FROM SPECIFIC CAUSES ARE PER 100,000 POPULATION, EXCEPT DEATHS FROM MATERNAL CAUSES ARE PER 100,000 LIVE BIRTHS AND DEATHS FROM CONDITIONS ORIG IN PERINATAL PERIOD ARE PER 1,000 LIVE BIRTHS. POPULATION FIGURES ARE 2003 ESTIMATES.

LESS THAN 0.06.

SELECTED DEATH STATISTICS, RESIDENTS OF MISSISSIPPI, 2003, continued

| AGE | TOTAL | | | MALE | | | FEMALE | | |
|----------------|--------|--------|-----------|--------|-------|-----------|--------|-------|-----------|
| | TOTAL | WHITE | NON-WHITE | TOTAL | WHITE | NON-WHITE | TOTAL | WHITE | NON-WHITE |
| TOTAL | 28,333 | 19,013 | 9,320 | 14,137 | 9,389 | 4,748 | 14,196 | 9,624 | 4,572 |
| UNDER 1 | 453 | 158 | 295 | 254 | 79 | 175 | 199 | 79 | 120 |
| (NEONATAL) | 246 | 78 | 168 | 134 | 40 | 94 | 112 | 38 | 74 |
| (POSTNEONATAL) | 207 | 80 | 127 | 120 | 39 | 81 | 87 | 41 | 46 |
| 1-4 | 88 | 38 | 50 | 48 | 21 | 27 | 40 | 17 | 23 |
| 5-9 | 44 | 22 | 22 | 22 | 13 | 9 | 22 | 9 | 13 |
| 10-14 | 65 | 35 | 30 | 37 | 20 | 17 | 28 | 15 | 13 |
| 15-24 | 479 | 260 | 219 | 326 | 171 | 155 | 153 | 89 | 64 |
| 25-34 | 655 | 302 | 353 | 430 | 205 | 225 | 225 | 97 | 128 |
| 35-44 | 1,260 | 677 | 583 | 734 | 401 | 333 | 526 | 276 | 250 |
| 45-54 | 2,400 | 1,276 | 1,124 | 1,508 | 847 | 661 | 892 | 429 | 463 |
| 55-64 | 3,504 | 2,236 | 1,268 | 2,127 | 1,387 | 740 | 1,377 | 849 | 528 |
| 65-74 | 5,184 | 3,535 | 1,649 | 2,936 | 2,062 | 874 | 2,248 | 1,473 | 775 |
| 75-84 | 7,453 | 5,570 | 1,883 | 3,582 | 2,711 | 871 | 3,871 | 2,859 | 1,012 |
| 85 AND OVER | 6,743 | 4,902 | 1,841 | 2,132 | 1,472 | 660 | 4,611 | 3,430 | 1,181 |
| UNKNOWN | 5 | 2 | 3 | 1 | 0 | 1 | 4 | 2 | 2 |

MORTALITY RATES**

| | | | |
|--------------|------|-----|------|
| INFANT | 10.7 | 6.8 | 15.4 |
| NEONATAL | 5.8 | 3.4 | 8.7 |
| POSTNEONATAL | 4.9 | 3.5 | 6.6 |

2003 POPULATION

| | |
|----------|-----------|
| TOTAL | 2,881,283 |
| WHITE | 1,763,609 |
| NONWHITE | 1,117,674 |

** DEATHS PER 1,000 LIVE BIRTHS.